


IN THE CLAIMS

1. (original): A process for the preparation of a compound of Formula (1):



Formula (1)


wherein:

R^x is optionally substituted aryl; and

R^y is optionally substituted hydrocarbyl;


which comprises the steps:

(a) reducing a compound of Formula (2):



Formula (2)


to a compound of Formula (3):



Formula (3)

wherein R^x and R^y are as defined for Formula (1):

(b) reacting a compound of Formula (3) with a leaving group donor, to give a compound of Formula (4);



Formula (4)


wherein:

R^x and R^y are as defined for Formula (1); and

OL is a leaving group:

(c) reacting a compound of Formula (4) with ammonia to give a compound of Formula (1).

2. (original): A process according to claim 1 for the preparation of a compound of Formula (5):



Formula (5)

wherein:


R^1 is a substituent;

R^2 is optionally substituted hydrocarbyl; and

n is 0 to 4:


which comprises the steps:

(a) reducing a compound of Formula (6):



Formula (6)


to a compound of Formula (7):



Formula (7)

wherein R^1 , R^2 and n are as defined for Formula (5):

(b) reacting a compound of Formula (7) with a leaving group donor, to give a compound of Formula (8);



Formula (8)


wherein:

R^1 , R^2 and n are as defined for Formula (5);

OL is a leaving group:

(c) reacting a compound of Formula (8) with ammonia to give a compound of Formula (5).

3. (original): A process according to claim 2 where R^2 is optionally substituted C_{1-4} alkyl.
4. (original): A process according to claim 3 where R^2 is methyl.
5. (currently amended): A process according to ~~any one of the preceding claims~~ claim 1 wherein n is 0.
6. (currently amended): A process according to ~~any one of the preceding claims~~ claim 1 where step (a) is carried out in the presence of a catalyst.
7. (original): A process according to claim 6 where the catalyst is of Formula (A):



Formula (A)

wherein:

R^3 represents a neutral optionally substituted hydrocarbyl, a neutral optionally substituted perhalogenated hydrocarbyl, or an optionally substituted cyclopentadienyl ligand;


A represents $-NR^4-$, $-NR^5-$, $-NHR^4$, $-NR^4R^5$ or $-NR^5R^6$ where R^4 is H, $C(O)R^6$, SO_2R^6 , $C(O)NR^6R^{10}$, $C(S)NR^6R^{10}$, $C(=NR^{10})SR^{11}$ or $C(=NR^{10})OR^{11}$, R^5 and R^6 each independently represents an optionally substituted hydrocarbyl, perhalogenated hydrocarbyl or an optionally substituted heterocyclyl group, and R^{10} and R^{11} are each independently hydrogen or a group as defined for R^6 ;

B represents $-O-$, $-OH$, OR^7 , $-S-$, $-SH$, SR^7 , $-NR^7-$, $-NR^8-$, $-NHR^8$, $-NR^7R^8$, $-NR^7R^9$, $-PR^7-$ or $-PR^7R^9$ where R^8 is H, $C(O)R^9$, SO_2R^9 , $C(O)NR^9R^{12}$, $C(S)NR^9R^{12}$, $C(=NR^{12})SR^{13}$ or $C(=NR^{12})OR^{13}$, R^7 and R^9 each independently represents an optionally substituted hydrocarbyl, perhalogenated hydrocarbyl or an optionally substituted heterocyclyl group, and R^{12} and R^{13} are each independently hydrogen or a group as defined for R^9 ;

E represents a linking group;

M represents a metal capable of catalysing transfer hydrogenation; and

COPY



Formula (9)

wherein:

R^1 is a substituent;


R^2 is optionally substituted hydrocarbyl; and

n is 0 to 4.

13. (currently amended): A process according to ~~any one of claims 1 to 5~~ claim 1 where in step (b) the leaving group donor is a compound of formula $R^{14}SO_2X$, where R^{14} is an optionally substituted alkyl, optionally substituted aryl or an optionally substituted heteroaryl group and X is a halogen.

14. (original): A process according to claim 13 where in step (b) the leaving group donor is methanesulphonyl chloride.


15. (currently amended): A process according to ~~either claim 1 or claim 2~~ for the preparation of a compound of Formula (10):



Formula (10)


which comprises the steps:

(a) reducing a compound of Formula (11):




Formula (11)

to a compound of Formula (12):



Formula (12)

(b) reacting a compound of Formula (12) with a compound of formula R^3SO_2X , in the presence of a base, to give a compound of Formula (13);



Formula (13)

wherein:

R^3 is optionally substituted C_{1-4} alkyl; and


X is halogen:

(c) reacting a compound of Formula (13) with ammonia to give a compound of Formula (10).

16. (original): A process according to claim 15 where step (a) is carried out in the presence of a catalyst of Formula (A) as described in claim 7.

17. (original): A process according to claim 15 wherein the compound of Formula (10) is purified by diastereomeric salt resolution using (L)-tartaric acid or (L)-chloropropionic acid.

18. (original): A process for the preparation of a stereoisomer of a compound of Formula (14):



Formula (14)


wherein:

R^1 is a substituent;

R^2 is optionally substituted hydrocarbyl; and

n is 0 to 4:

which comprises the transfer hydrogenation of a compound of Formula (6):




Formula (6)

by a hydrogen donor in the presence of a catalyst of Formula (A) as described in claim 7.

19. (original): A process for the diastereomeric salt resolution of (S)-1-naphthylethylamine which comprises mixing (S)-1-naphthylethylamine with (2R,3R)-tartaric acid or (S)-chloropropionic acid to form the corresponding diastereomeric salt.

20. (original): A diastereomeric salt of (S)-1-naphthylethylamine with (2R,3R)-tartaric acid or (S)-chloropropionic acid.

21. (original): A compound of Formula (15):



Formula (15)


wherein:

R^1 is a substituent;

R^2 is optionally substituted hydrocarbyl; and

n is 0 to 4.

22. (original): A compound according to claim 21 of Formula (15) which is of Formula (16):



Formula (16)